REFERRAL GUIDELINES - BREAST CLINIC



Please review guidelines before referring patients. Please include as much detail as possible to assist us to accurately triage the referral. Fax referrals to 8345 6856.

Specialist review prior to image guided or clinician biopsy of lesions is preferred.

Where imaging is undertaken, please advise patients to

- Take previous films for the radiologist to review AND
- Bring films/disks to their hospital appointment.

Family history guidelines:

Specify all family members including whether they are a 1st or 2nd degree relative, their relationship to the patient and age of diagnosis if known who have had:

• breast cancer +/or ovarian cancer, bilateral cancer or male breast cancer.

Please include whether the patient is of Ashkenazi heritage.

SYMPTOMS	RECOMMENDATION	
Breast Pain		
Chest wall pain	GP to treat	
Thickening or nodularity consistent with hormonal change	Review in 2 or 3 months	
Discrete lesion	Yes – Imaging* then refer	No - Review in 2 to 3 months
Pain/nodularity persists after 2 months	Imaging* then refer	
Lumps		
Thickening or nodularity with hormonal change	Clinical review in 2 or 3 months	
Discrete lump(s)	Imaging* then refer	
Nipple Discharge and Change		
Bilateral on expression only	Reassure, encourage not to express.	
No discrete lesion	Review in 2-3 months	
No blood		
Unilateral spontaneous discharge	Imaging* then refer	
Blood discharge	Imaging* then refer	
Single duct discharge		
Nipple inversion/ change (< 3 months)		
Nipple eczema (i.e. new)	Imaging* and refer	

RESULTS OF IMAGING	RECOMMENDATION	
Normal or No discrete lesion (Category 1)		
Consistent with Clinical exam	Reassure patient. Advise Breast self awareness Advise age >50 two yearly mammogram – BreastScreen 13 20 50 http://www.breastscreen.org.au/	
Inconsistent	Refer	
Benign (Category 2)		
Simple cysts	Not palpable - reassure	
	Palpable – refer for aspiration	
Complex cysts	Not palpable: • Age <40 years old - reassure and re-examine 6/12 • Age >40 years old - re-examine and re image 6/12 Palpable - refer	
Solid lesions (plus no clinical abnormality) Age < 25 years old	Reassure	
Age 25 – 35 years old	Repeat ultrasound at 6 months + stop if no change	
Age > 35 years old	Refer	
Indeterminate (Category 3) and Suspicious (Category 4) / Malignant		
	Refer as per Breast Cancer Referral Guidelines	