Reproductive medicine – A new Reproductive Choices service offering surgical termination of pregnancy between 7 – 13+6 gestation for pregnancies for disadvantaged women with strict limited eligibility criteria is commencing in November 2023. This service does not offer medical termination of pregnancy except for miscarriages.GP referral required with relevant dating ultrasound and bloods.

Reproductive Choices ServicesDisadvantaged women:Dating UltrasoundEligibility criteria aligns with a strategy of equity of access for vulnerable populations. For• Dating Ultrasound	Primary Condition	Key Information Points	Clinical Investigation
example: • Aboriginal or Torres Strait Islander • Addiction or substance abuse • Adolescent (age <25) • Family Violence • Healthcare card holders • History of incarceration • Patient age >/= 14 of age referred to the Royal Women's Hospital • No fixed address • Refugee background • Gestational Age (GA) – up to 13 weeks + 6 days • Significant mental health/ psychosocial issues • Speak English as a second Language	Reproductive Choices Services	 Eligibility criteria aligns with a strategy of equity of access for vulnerable populations. For example: Aboriginal or Torres Strait Islander Addiction or substance abuse Adolescent (age <25) Family Violence Healthcare card holders History of incarceration Patient age >/= 14 of age referred to the Royal Women's Hospital No fixed address Refugee background Gestational Age (GA) – up to 13 weeks + 6 days Significant mental health/psychosocial issues Speak English as a second 	Relevant bloods -BCHG,

Access and referral priority Gynaecology

The clinical information provided in your referral will determine the triage category. The triage category will affect the timeframe in which the patient is offered an appointment.

URGENT	URGENT	ROUTINE
Appointment timeframe 30 days.	Appointment within 60	Appointment timeframe greater
	days	than 30 days, depending on clinical need.
 Post-menopausal bleeding, including any pre malignant conditions including VIN/VAIN/ complex endometrial hyperplasia EPAS (Early Pregnancy). Abnormal uterine bleeding with Hb < 100g/DL or severe quality of life impairment. Adnexal (ovary/tube) abnormality assessed as at risk of malignancy, torsion or other significant complication. Other pelvic mass/tumour assessed as significant risk of malignancy or with severe symptoms. (e.g. fibroids with recent increase in size or pain) Pelvic organ prolapse with urinary retention or Quality Of Life change assessed as severe and disabling. Urinary retention or Voiding dysfunction emptying ≤50% of bladder volume. Undiagnosed pelvic pain requiring hospital management. Vulval conditions with suspected malignancy. Persistent or recurring post-coital bleeding Disadvantaged Women requesting Surgical Termination of Pregnancy 7-13+6 weeks gestation) 	 All colposcopy referrals are managed as urgent to be seen within 8 weeks as per National Cervical Screening Guidelines. Note: current capacity does not allow this to be achieved – all colposcopy referrals are assessed and prioritised according to risk) All OASIS (complex perineal and anal sphincter injuries 3B or more) are seen within 6-12 weeks of postnatal discharge. Pelvic floor symptoms or complications following incontinence or pelvic organ prolapse mesh surgery See: http://tiny.cc/whmms 	 Fertility referrals Contraception – reversible or permanent Menopause management Abnormal uterine bleeding not meeting criteria for 'URGENT' referral. Adnexal abnormalities with low risk of malignancy or other complication, particularly incidental findings on imaging. Uterine fibroids with minimal or no symptoms. Vulval conditions without risk of malignancy Persistent or chronic pelvic pain Pelvic organ prolapse not meeting 'URGENT' criteria. Lower urinary tract symptoms including urgency, frequency, incontinence (will be triaged for physiotherapy assessment prior to appointment), recurrent UTIs or bladder pain Voiding dysfunction not meeting 'URGENT' criteria Isolated haematuria (refer to Urology) Sexual dysfunction Labial surgery for medical indications.